

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## MISCELLANEOUS.

110. Proposed by E. W. MORRELL, South Tunbridge, Vt.

If a and b be the sides of a triangle, A and B the angles opposite, then will  $\log b - \log a = \cos 2A - \cos 2B + \frac{1}{2}(\cos 4A - \cos 4B) + \frac{1}{8}(\cos 6A - \cos 6B) + \dots$ 

111. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Exhibit  $\cos^3\theta \sin^3\theta \sin^2\phi \cos\phi$  as a series of harmonics.

- 112. Proposed by LON C. WALKER, A. M., Assistant Professor of Mathematics, Leland Stanford Jr. University, Palo Alto, Cal.
- (a) Find the area enclosed by four circles, of which two touch the x-axis, and two the y-axis, at the origin.
- (b) Required the area enclosed by four parabolas, of which two touch the x-axis, and two the y-axis, at the origin.
- 113. Proposed by F. P. MATZ. M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pa.

Deduce the Sylvestrian Reciprocant from  $x^4+y^4=4x^2y^2$ .

\*\* Solutions of these problems should be sent to J. M. Colaw not later than August 10.

## NOTES.

Dr. W. B. Fite has been appointed Instructor in Mathematics at Cornell University. He will take most of the advanced work formerly done by Dr. G. A. Miller.

Professor Alexander S. Chessin has been elected Professor of Mathematics in the Washington University, St. Louis. The University is to be congratulated in securing such an able man.

Mr. Paul A. Towne, A. M., has sent us a number of very interesting magic squares constructed on the sides and hypothenuse of right triangles. but owing to the difficulty in reproducing them, we must omit their publication.

Brief notice will be made in the next issue of the Monthly of the following books: One Hundred Problems in Mathematical Physics, by E. P. Thompson; Atoms and Energy, by D. A. Murray; Experimental Physics, by Eugene Lommel; An Introduction to the Study of Chemistry, by Ira Remsen.